

MAS151: Civil Engineering Mathematics

Dr James Cranch

mas-engineering@sheffield.ac.uk

Tuesday 31st March 2020, 12pm
Hicks LT1

End of year exam

The exam for this course will happen in the May/June exam period.

The exam for this course will happen in the May/June exam period. The draft timetable has not been announced yet

The exam for this course will happen in the May/June exam period. The draft timetable has not been announced yet so please check the final timetable!

The exam for this course will happen in the May/June exam period. The draft timetable has not been announced yet so please check the final timetable!

The exam will count for 85% of your final grade (the other 15% comes from online tests).

The exam for this course will happen in the May/June exam period. The draft timetable has not been announced yet so please check the final timetable!

The exam will count for 85% of your final grade (the other 15% comes from online tests).

You could be examined on anything from the module.

The exam for this course will happen in the May/June exam period. The draft timetable has not been announced yet so please check the final timetable!

The exam will count for 85% of your final grade (the other 15% comes from online tests).

You could be examined on anything from the module.

Here are some details about the exam, and tips for how to do well on maths exams.

Exam format

The exam will be 3 hours long.

The exam will be 3 hours long. It will consist of

The exam will be 3 hours long. It will consist of

- Section A: 12 compulsory questions, 3 marks each;

The exam will be 3 hours long. It will consist of

- Section A: 12 compulsory questions, 3 marks each;
- Section B: 8 compulsory questions, 8 marks each.

The exam will be 3 hours long. It will consist of

- Section A: 12 compulsory questions, 3 marks each;
- Section B: 8 compulsory questions, 8 marks each.

Section A questions are intended to be straightforward tests of skills.

The exam will be 3 hours long. It will consist of

- Section A: 12 compulsory questions, 3 marks each;
- Section B: 8 compulsory questions, 8 marks each.

Section A questions are intended to be straightforward tests of skills. Section B questions will require a bit more thought.

The exam will be 3 hours long. It will consist of

- Section A: 12 compulsory questions, 3 marks each;
- Section B: 8 compulsory questions, 8 marks each.

Section A questions are intended to be straightforward tests of skills. Section B questions will require a bit more thought.

To pass the module comfortably, you will need to do well on Section B

The exam will be 3 hours long. It will consist of

- Section A: 12 compulsory questions, 3 marks each;
- Section B: 8 compulsory questions, 8 marks each.

Section A questions are intended to be straightforward tests of skills. Section B questions will require a bit more thought.

To pass the module comfortably, you will need to do well on Section B as Section A only carries 36 marks.

Past papers and formula sheet

There are two past papers on the course webpage with solutions.

There are two past papers on the course webpage with solutions.

You can find exams for earlier years on the School of Mathematics and Statistics webpage

There are two past papers on the course webpage with solutions.

You can find exams for earlier years on the School of Mathematics and Statistics webpage (MAS140 has the earliest papers)

There are two past papers on the course webpage with solutions.

You can find exams for earlier years on the School of Mathematics and Statistics webpage (MAS140 has the earliest papers) but we will not provide solutions to them.

There are two past papers on the course webpage with solutions.

You can find exams for earlier years on the School of Mathematics and Statistics webpage (MAS140 has the earliest papers) but we will not provide solutions to them.

Also on the course webpage is the formula sheet which will be provided to you in the exam.

There are two past papers on the course webpage with solutions.

You can find exams for earlier years on the School of Mathematics and Statistics webpage (MAS140 has the earliest papers) but we will not provide solutions to them.

Also on the course webpage is the formula sheet which will be provided to you in the exam. You can use any results from this formula sheet without proof.

Calculators

Remember that you need to have an approved calculator for use in the exam.

Remember that you need to have an approved calculator for use in the exam. There is a page on the University web page which gives full information.

Remember that you need to have an approved calculator for use in the exam. There is a page on the University web page which gives full information.

www.shef.ac.uk/ssid/exams/calculator

Remember that you need to have an approved calculator for use in the exam. There is a page on the University web page which gives full information.

www.shef.ac.uk/ssid/exams/calculator

You will need a calculator for the exam!

Remember that you need to have an approved calculator for use in the exam. There is a page on the University web page which gives full information.

www.shef.ac.uk/ssid/exams/calculator

You will need a calculator for the exam!

Some approved calculators have functions that could allow you to provide answers without doing the calculations yourself.

Remember that you need to have an approved calculator for use in the exam. There is a page on the University web page which gives full information.

www.shef.ac.uk/ssid/exams/calculator

You will need a calculator for the exam!

Some approved calculators have functions that could allow you to provide answers without doing the calculations yourself.

But please note that we expect to see fully worked solutions to all questions.

Unfair means

You must not bring in anything to the exam which breaks the University's exam regulations.

You must not bring in anything to the exam which breaks the University's exam regulations.

In particular, you must not bring in any notes or revision materials,

You must not bring in anything to the exam which breaks the University's exam regulations.

In particular, you must not bring in any notes or revision materials, and there must be no hidden writing on your pencil case or calculator.

You must not bring in anything to the exam which breaks the University's exam regulations.

In particular, you must not bring in any notes or revision materials, and there must be no hidden writing on your pencil case or calculator.

People do get caught cheating, and the consequences can be severe!

You must not bring in anything to the exam which breaks the University's exam regulations.

In particular, you must not bring in any notes or revision materials, and there must be no hidden writing on your pencil case or calculator.

People do get caught cheating, and the consequences can be severe!

There is a video about the exam process on the Exams Office page on the University website.

You must not bring in anything to the exam which breaks the University's exam regulations.

In particular, you must not bring in any notes or revision materials, and there must be no hidden writing on your pencil case or calculator.

People do get caught cheating, and the consequences can be severe!

There is a video about the exam process on the Exams Office page on the University website.

www.shef.ac.uk/ssid/exams

Preparing for maths exams

Practice

To become good at mathematical techniques requires lots of practice.

To become good at mathematical techniques requires lots of practice.

The examples you have worked on in the problem classes should have deepened your understanding of the maths,

To become good at mathematical techniques requires lots of practice.

The examples you have worked on in the problem classes should have deepened your understanding of the maths, but to do well on the exam we strongly recommend that you work through the exercise sheets on the course webpage.

To become good at mathematical techniques requires lots of practice.

The examples you have worked on in the problem classes should have deepened your understanding of the maths, but to do well on the exam we strongly recommend that you work through the exercise sheets on the course webpage.

If you have questions on the problems on the exercise sheets, feel free to ask in your problem class or post on the discussion board.

Past papers (again)

I would expect everyone to practice on the past papers before the exam.

I would expect everyone to practice on the past papers before the exam. It can be good to do these under timed conditions so that you know whether you will be under time pressure or not.

I would expect everyone to practice on the past papers before the exam. It can be good to do these under timed conditions so that you know whether you will be under time pressure or not.

However, you should not expect the questions to be identical to the ones from last year!

Presentation

In Semester 1, I tried to convince you about the importance of presenting work well.

In Semester 1, I tried to convince you about the importance of presenting work well. It's hard to present work well in exams,

In Semester 1, I tried to convince you about the importance of presenting work well. It's hard to present work well in exams, but it's probably the time when it's most important to make your working clear.

In Semester 1, I tried to convince you about the importance of presenting work well. It's hard to present work well in exams, but it's probably the time when it's most important to make your working clear. As a reminder,

In Semester 1, I tried to convince you about the importance of presenting work well. It's hard to present work well in exams, but it's probably the time when it's most important to make your working clear. As a reminder,

- use words to explain what you are doing;

In Semester 1, I tried to convince you about the importance of presenting work well. It's hard to present work well in exams, but it's probably the time when it's most important to make your working clear. As a reminder,

- use words to explain what you are doing;
- link together equal expressions with $=$ -signs, and line them up if possible;

In Semester 1, I tried to convince you about the importance of presenting work well. It's hard to present work well in exams, but it's probably the time when it's most important to make your working clear. As a reminder,

- use words to explain what you are doing;
- link together equal expressions with $=$ -signs, and line them up if possible;
- avoid floating answers;

In Semester 1, I tried to convince you about the importance of presenting work well. It's hard to present work well in exams, but it's probably the time when it's most important to make your working clear. As a reminder,

- use words to explain what you are doing;
- link together equal expressions with $=$ -signs, and line them up if possible;
- avoid floating answers;
- avoid backwards reasoning;

In Semester 1, I tried to convince you about the importance of presenting work well. It's hard to present work well in exams, but it's probably the time when it's most important to make your working clear. As a reminder,

- use words to explain what you are doing;
- link together equal expressions with $=$ -signs, and line them up if possible;
- avoid floating answers;
- avoid backwards reasoning;
- write concluding statements.

In Semester 1, I tried to convince you about the importance of presenting work well. It's hard to present work well in exams, but it's probably the time when it's most important to make your working clear. As a reminder,

- use words to explain what you are doing;
- link together equal expressions with $=$ -signs, and line them up if possible;
- avoid floating answers;
- avoid backwards reasoning;
- write concluding statements.

In Semester 1, I tried to convince you about the importance of presenting work well. It's hard to present work well in exams, but it's probably the time when it's most important to make your working clear. As a reminder,

- use words to explain what you are doing;
- link together equal expressions with $=$ -signs, and line them up if possible;
- avoid floating answers;
- avoid backwards reasoning;
- write concluding statements.

Make sure all the steps in your calculations are clearly presented.

Activity. Look at the two sets of genuine student solutions to questions on material from this course. In pairs, discuss what score (out of 10) you think they got.

I'll reveal the questions and the marks shortly.

The first question is

The first question is

- (i) Give the first 4 terms in the binomial expansions of
(a) $(1 - 3x)^5$; (b) $(8 + x)^{\frac{1}{3}}$. (7 marks)

The first question is

- (i) Give the first 4 terms in the binomial expansions of
(a) $(1 - 3x)^5$; (b) $(8 + x)^{\frac{1}{3}}$. (7 marks)
- (ii) By setting $x = 0.1$ in your answer to (b), find an approximate value for the cube root of 8.1, accurate to 4 decimal places. (3 marks)

The first question is

- (i) Give the first 4 terms in the binomial expansions of
(a) $(1 - 3x)^5$; (b) $(8 + x)^{\frac{1}{3}}$. (7 marks)
- (ii) By setting $x = 0.1$ in your answer to (b), find an approximate value for the cube root of 8.1, accurate to 4 decimal places. (3 marks)

Maths Coursework

(A)

i) $a) (1-3x)^5 = 1 + (-3x) + (-3x)^2 + (-3x)^3$

$$= 1 - 3x + 9x^2 - 27x^3$$

b) $(8+x)^{1/3} = 8^{1/3} (1 + \frac{1}{8}x)^{1/3} = 2 (1 + \frac{1}{8}x)^{1/3}$

~~$$= 2(1 + \frac{1}{8}x)^{1/3}$$~~

$$= 2 \left(1 + \frac{1}{8} \left(\frac{1}{8}x \right) + \frac{\binom{1/3}{2} \left(\frac{1}{8}x \right)^2}{2} + \frac{\binom{1/3}{3} \left(\frac{1}{8}x \right)^3}{6} \right)$$

$$= 2 \left(1 + \frac{1}{24}x - \frac{2}{9} \left(\frac{1}{64}x^2 \right) + \frac{10}{27} \left(\frac{1}{512}x^3 \right) \right)$$

$$= 2 + \frac{1}{12}x - \frac{1}{9} \left(\frac{1}{64}x^2 \right) + \frac{5}{27} \left(\frac{1}{512}x^3 \right)$$

$$= 2 + \frac{1}{12}x - \frac{1}{576}x^2 + \frac{5}{41472}x^3$$

ii) $x=0.1$

$$2 + \frac{1}{12}(0.1) - \frac{1}{576}(0.1)^2 + \frac{5}{41472}(0.1)^3$$

$$= 2 + \frac{1}{120} - 1.7361 \times 10^{-5} + 1.2056 \times 10^{-7}$$

$$= 2.0083$$

① (i) $a) (1-3x)^5 = 1 + \frac{5(-3x)}{1!} + \frac{(5)(4)(-3x)^2}{2!} + \frac{(5)(4)(3)(-3x)^3}{3!}$

$$= 1 - 15x + \frac{(20)(9x^2)}{2} - \frac{(60)(27x^3)}{6}$$

Answer = $1 - 15x + 90x^2 - 270x^3$

(B)

b) $(8+x)^{1/3}$
 $= 8^{1/3} \left(1 + \frac{x}{8} \right)^{1/3}$
 $= 2 \left[1 + \frac{\binom{1/3}{1} \left(\frac{1}{8}x \right)}{1!} + \frac{\binom{1/3}{2} \left(\frac{1}{8}x \right)^2}{2!} + \frac{\binom{1/3}{3} \left(\frac{1}{8}x \right)^3}{3!} \right]$

$$= 2 \left[1 + \frac{\frac{1}{24}x}{1} + \frac{\left(-\frac{1}{2} \right) \left(\frac{1}{64}x^2 \right)}{2} + \frac{\left(\frac{10}{27} \right) \left(\frac{1}{512}x^3 \right)}{6} \right]$$

$$= 2 \left[1 + \frac{1}{24}x - \frac{1}{576}x^2 + \frac{5}{41472}x^3 \right]$$

Answer = $2 + \frac{1}{12}x - \frac{1}{288}x^2 + \frac{5}{20736}x^3$

(ii) $2 + \left(\frac{1}{2} \right) (0.1) - \left(\frac{1}{288} \right) (0.1)^2 + \left(\frac{5}{20736} \right) (0.1)^3$
 $= 2 + \frac{1}{20} - \frac{1}{28800} + \frac{1}{4147200}$
 $= 2.008248852$

Answer = 2.0083 (4.d.p.)

E

i) $a) (1-3x)^5 = 1 + (-3x) + \frac{(-3x)^2}{2!} + \frac{(-3x)^3}{3!}$

$$1 - 3x + 3x^2 - 30x^3$$

1) $b) (8+x)^{1/3} = 8^{1/3} \left(1 + \frac{1}{8}x \right)^{1/3} = 2 \left[1 + \frac{1}{8}x + \frac{\binom{1/3}{2} \left(\frac{1}{8}x \right)^2}{2!} + \frac{\binom{1/3}{3} \left(\frac{1}{8}x \right)^3}{3!} \right]$

$$2 + \frac{1}{4}x - \frac{1}{36}x^2 + \frac{5}{324}x^3$$

ii) $x=0.1$ $2 + \frac{1}{4}(0.1) - \frac{1}{36}(0.1)^2 + \frac{5}{324}(0.1)^3 = 2.00804$

Answer = 2.0080 (4.d.p.)

The students scored

The students scored

A: 6,

The students scored

A: 6, B: 10,

The students scored

A: 6, B: 10, E: 0.

Maths Coursework

(A)

d) i) a) $(1-3x)^5 = 1 + (-3x) + (-3x)^2 + (-3x)^3$

$$= 1 - 3x + 9x^2 - 27x^3$$

b) $(8+x)^{\frac{1}{3}} = 8^{\frac{1}{3}} (1 + \frac{1}{8}x)^{\frac{1}{3}} = 2(1 + \frac{1}{8}x)^{\frac{1}{3}}$

~~$$= 2(1 + \frac{1}{8}x)^{\frac{1}{3}}$$~~

$$= 2 \left(1 + \frac{1}{3}(\frac{1}{8}x) + \frac{(\frac{1}{3})(\frac{1}{3})(\frac{1}{8}x)^2}{2} + \frac{(\frac{1}{3})(\frac{1}{3})(\frac{1}{3})(\frac{1}{8}x)^3}{6} \right)$$

$$= 2 \left(1 + \frac{1}{24}x + \frac{-2}{9} \left(\frac{1}{64}x^2 \right) + \frac{10}{27} \left(\frac{1}{512}x^3 \right) \right)$$

$$= 2 + \frac{1}{12}x - \frac{1}{9} \left(\frac{1}{64}x^2 \right) + \frac{5}{27} \left(\frac{1}{512}x^3 \right)$$

$$= 2 + \frac{1}{12}x - \frac{1}{576}x^2 + \frac{5}{41472}x^3$$

ii) $x=0.1$

$$2 + \frac{1}{12}(0.1) - \frac{1}{576}(0.1)^2 + \frac{5}{41472}(0.1)^3$$

$$= 2 + \frac{1}{120} - 1.7361 \times 10^{-5} + 1.2056 \times 10^{-7}$$

$$= 2.0083$$

① (i) a) $(1-3x)^5 = 1 + \frac{5(-3x)}{1!} + \frac{(5)(4)(-3x)^2}{2!} + \frac{(5)(4)(3)(-3x)^3}{3!}$

$$= 1 - 15x + \frac{(20)(9x^2)}{2} - \frac{(60)(27x^3)}{6}$$

Answer = $1 - 15x + 90x^2 - 270x^3$

(b) $(8+x)^{\frac{1}{3}}$

$$= 8^{\frac{1}{3}} \left(1 + \frac{x}{8} \right)^{\frac{1}{3}}$$

$$= 2 \left[1 + \frac{(\frac{1}{3})(\frac{1}{8}x)}{1!} + \frac{(\frac{1}{3})(\frac{1}{3})(\frac{1}{8}x)^2}{2!} + \frac{(\frac{1}{3})(\frac{1}{3})(\frac{1}{3})(\frac{1}{8}x)^3}{3!} \right]$$

$$= 2 \left[1 + \frac{\frac{1}{24}x}{1} + \frac{(-\frac{2}{9})(\frac{1}{64}x^2)}{2} + \frac{(\frac{10}{27})(\frac{1}{512}x^3)}{6} \right]$$

$$= 2 \left[1 + \frac{1}{24}x - \frac{1}{576}x^2 + \frac{5}{41472}x^3 \right]$$

Answer = $2 + \frac{1}{12}x - \frac{1}{288}x^2 + \frac{5}{20736}x^3$

(ii) $2 + \left(\frac{1}{12}\right)(0.1) - \left(\frac{1}{288}\right)(0.1)^2 + \left(\frac{5}{20736}\right)(0.1)^3$

$$= 2 + \frac{1}{120} - \frac{1}{28800} + \frac{5}{4147200}$$

$$= 2.008248852$$

Answer = 2.0083 (4 d.p.)

E

i) a) $(1-3x)^5 = 1 + (-3x) + \frac{(-3x)^2}{2!} + \frac{(-3x)^3}{3!}$

$$1 - 3x + 3x^2 - 30x^3$$

1) b) $(8+x)^{\frac{1}{3}} = 8^{\frac{1}{3}} \left(1 + \frac{1}{8}x \right)^{\frac{1}{3}} = 2 \left[1 + \frac{1}{24}x - \frac{1}{576}x^2 + \frac{5}{41472}x^3 \right]$

$$2 + \frac{1}{12}x - \frac{1}{288}x^2 + \frac{5}{20736}x^3$$

ii) $x=0.1$ $2 + \frac{1}{12}(0.1) - \frac{1}{288}(0.1)^2 + \frac{5}{20736}(0.1)^3 = 2.00804$

Answer = 2.0080 (4 d.p.)

The second question is

The second question is

- (i) Obtain the first three nonzero terms of the Taylor series expansion of $f(x) = \cosh(x - 2)$ around $x = 2$. (6 marks)

The second question is

- (i) Obtain the first three nonzero terms of the Taylor series expansion of $f(x) = \cosh(x - 2)$ around $x = 2$. (6 marks)
- (ii) Using the expansion obtained in (i), find the two approximate solutions to the equation $\cosh(x - 2) = 1.1$. [Hint: Only the solution of a quadratic equation is required]. (4 marks)

The second question is

- (i) Obtain the first three nonzero terms of the Taylor series expansion of $f(x) = \cosh(x - 2)$ around $x = 2$. (6 marks)
- (ii) Using the expansion obtained in (i), find the two approximate solutions to the equation $\cosh(x - 2) = 1.1$. [Hint: Only the solution of a quadratic equation is required]. (4 marks)

C 3) i) $f(x) = \cosh(x-2)$ $a=2$

(A)

$$f(a) + f'(a)(x-a) + \frac{f''(a)}{2!}(x-a)^2$$

$$= \cosh(2-2) + \sinh(2-2)(x-2) + \cosh(0)(x-2)^2$$

$$+ \frac{\cosh(0)}{2!}(x-2)^2 + \frac{\cosh(0)}{4!}(x-2)^4$$

$$= 1 + (x-2)^2 + (x-2)^4$$

ii) $1 + (x-2)^2 + (x-2)^4 = 1$

$$\cosh(x-2) = 1.1$$

$$1 + (x-2)^2 = 1.1$$

$$(x-2)^2 = 0.1 \quad x-2 = \pm \sqrt{0.1}$$

$$x = 2 \pm \sqrt{0.1}$$

$$x^2 - 4x + 3.6 = 0$$

$$(x-2)^2 - 0.4 = 0$$

$$(x-2)^2 = 0.4 \quad x-2 = \pm \sqrt{0.4}$$

③ (i) $f(x) = \cosh(x-2)$ around $x=2$ $f(0.12) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)x^n}{n!}$

$$f'(x) = \sinh(x-2) \quad f'(2) = \sinh(0) = 0$$

$$f''(x) = \cosh(x-2) \quad f''(2) = \cosh(0) = 1$$

$$f'''(x) = \sinh(x-2) \quad f'''(2) = \sinh(0) = 0$$

$$f^{(4)}(x) = \cosh(x-2) \quad f^{(4)}(2) = \cosh(0) = 1$$

(B)

$$f(x) = 1 + \frac{0(x-2)}{1!} + \frac{1(x-2)^2}{2!} + \frac{0(x-2)^3}{3!} + \frac{1(x-2)^4}{4!}$$

$$\text{Answer: } 1 + \frac{(x-2)^2}{2} + \frac{(x-2)^4}{24}$$

(ii) $\cosh(x-2) = 1.1$

$$\therefore 1 + \frac{(x-2)^2}{2} + \frac{(x-2)^4}{24} = 1.1$$

$$24 + 12(x-2)^2 + (x-2)^4 = 26.4$$

$$(x-2)^4 + 12(x-2)^2 - 2.4 = 0$$

$$\text{let } y = x-2$$

$$y^4 + 12y^2 - 2.4 = 0$$

$$\text{let } y^2 = z$$

$$z^2 + 12z - 2.4 = 0$$

$$\text{then use quadratic formula: } z = \frac{-30 \pm \sqrt{900}}{5} \quad z = \frac{-30 \pm 30}{5}$$

must be positive as $z = y^2$

$$y = \sqrt{z} = \sqrt{\frac{-30 + 30\sqrt{5}}{5}} = \pm 0.4435914268$$

$$x = y + 2$$

$$x = 2.443591427 \quad \text{and} \quad 1.556408573$$

$$\text{Solutions are: } 2.4436 \quad \text{and} \quad 1.5564 \quad (\text{4 d.p.})$$

The students scored

The students scored

A: 4,

The students scored

A: 4, B: 10.

3) i) $f(x) = \cosh(x-2)$ $a=2$

(A)

$$f(a) + f'(a)(x-a) + \frac{f''(a)}{2!}(x-a)^2$$

$$= \cosh(2-2) + \sinh(2-2)(x-2) + \cosh(0)(x-2)^2$$

$$+ \frac{\cosh(0)}{2!}(x-2)^2 + \frac{\cosh(0)}{4!}(x-2)^4$$

$$= 1 + (x-2)^2 + (x-2)^4$$

ii) $1 + (x-2)^2 + (x-2)^4 = 1$

$$\cosh(x-2) = 1.1$$

$$1 + (x-2)^2 = 1.1$$

$$(x-2)^2 = 0.1 \quad x-2 = \pm \sqrt{0.1}$$

$$x = 2 \pm \sqrt{0.1}$$

$$x^2 - 4x + 3.6 = 0$$

$$(x-2)^2 - 0.4 = 0$$

$$(x-2)^2 = 0.4 \quad x-2 = \pm \sqrt{0.4}$$

3) (i) $f(x) = \cosh(x-2)$ around $x=2$ $f(a+x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)x^n}{n!}$

$$f'(x) = \sinh(x-2) \quad f'(2) = \sinh(0) = 0$$

$$f''(x) = \cosh(x-2) \quad f''(2) = \cosh(0) = 1$$

$$f'''(x) = \sinh(x-2) \quad f'''(2) = \sinh(0) = 0$$

$$f^{(4)}(x) = \cosh(x-2) \quad f^{(4)}(2) = \cosh(0) = 1$$

(B)

$$f(x) = 1 + \frac{0(x-2)}{1!} + \frac{1(x-2)^2}{2!} + \frac{0(x-2)^3}{3!} + \frac{1(x-2)^4}{4!}$$

$$\text{Answer: } 1 + \frac{(x-2)^2}{2} + \frac{(x-2)^4}{24}$$

(ii) $\cosh(x-2) = 1.1$

$$\therefore 1 + \frac{(x-2)^2}{2} + \frac{(x-2)^4}{24} = 1.1$$

$$24 + 12(x-2)^2 + (x-2)^4 = 26.4$$

$$(x-2)^4 + 12(x-2)^2 - 2.4 = 0$$

$$\text{let } y = x-2$$

$$y^4 + 12y^2 - 2.4 = 0$$

$$\text{let } y^2 = z$$

$$z^2 + 12z - 2.4 = 0$$

$$\text{then use quadratic formula: } z = \frac{-30 \pm \sqrt{900}}{5} \quad z = \frac{-30 \pm 30}{5}$$

must be positive as $z = y^2$

$$y = \sqrt{z} = \sqrt{\frac{-30 + 30\sqrt{5}}{5}} = \pm 0.4435914268$$

$$x = y + 2$$

$$x = 2.443591427 \quad \text{and} \quad 1.556408573$$

$$\text{Solutions are: } 2.4436 \quad \text{and} \quad 1.5564 \quad (\text{4 d.p.})$$

Mathematical exam questions

There are some helpful things to bear in mind when doing maths exams.

Linked parts

The two questions we looked at both followed a similar format:

The two questions we looked at both followed a similar format: the first part of the question helped with the second part.

The two questions we looked at both followed a similar format: the first part of the question helped with the second part. In these examples, you were told to use the previous part.

The two questions we looked at both followed a similar format: the first part of the question helped with the second part.

In these examples, you were told to use the previous part.

Sometimes you won't be explicitly told to use your previous working, but it is expected you would spot that it's useful.

The two questions we looked at both followed a similar format: the first part of the question helped with the second part.

In these examples, you were told to use the previous part.

Sometimes you won't be explicitly told to use your previous working, but it is expected you would spot that it's useful.

Always think about whether the parts to a question are linked.

The two questions we looked at both followed a similar format: the first part of the question helped with the second part.

In these examples, you were told to use the previous part.

Sometimes you won't be explicitly told to use your previous working, but it is expected you would spot that it's useful.

Always think about whether the parts to a question are linked. Often you will find they are.

Hints

You may sometimes be given an obvious hint

You may sometimes be given an obvious hint (e.g. 'Hint: Use the binomial theorem').

You may sometimes be given an obvious hint (e.g. 'Hint: Use the binomial theorem').

More likely, it will be phrased as

You may sometimes be given an obvious hint (e.g. 'Hint: Use the binomial theorem').

More likely, it will be phrased as 'By using the binomial theorem, show that ...'.

You may sometimes be given an obvious hint (e.g. 'Hint: Use the binomial theorem').

More likely, it will be phrased as 'By using the binomial theorem, show that ...'. You may not always see how to use the hint straight away, but it will always help.

You may sometimes be given an obvious hint (e.g. 'Hint: Use the binomial theorem').

More likely, it will be phrased as 'By using the binomial theorem, show that ...'. You may not always see how to use the hint straight away, but it will always help.

A variation of 'By using the binomial theorem, or otherwise, show that ...' usually indicates there are other ways to proceed,

You may sometimes be given an obvious hint (e.g. 'Hint: Use the binomial theorem').

More likely, it will be phrased as 'By using the binomial theorem, show that ...'. You may not always see how to use the hint straight away, but it will always help.

A variation of 'By using the binomial theorem, or otherwise, show that ...' usually indicates there are other ways to proceed, but again the hint will normally help. But be aware that sometimes other methods can be easier.

Another phrase that is sometimes used is 'Hence show that...'.

Another phrase that is sometimes used is 'Hence show that...'. This means 'from your previous work, show that...'.

Another phrase that is sometimes used is 'Hence show that...'. This means 'from your previous work, show that...'.

Again, this sometimes comes in the form 'Hence, or otherwise, show that...'.

Another phrase that is sometimes used is 'Hence show that...'. This means 'from your previous work, show that...'.

Again, this sometimes comes in the form 'Hence, or otherwise, show that...'.

Look out for these kind of phrases, as they are there to help!

Understanding what's required

Sometimes questions are more straight-forward than people realise.

Sometimes questions are more straight-forward than people realise. For example,

Show that the function $y = a \cos x + b \sin x$ is a solution to

$$\frac{d^2y}{dx^2} + y = 0$$

Sometimes questions are more straight-forward than people realise. For example,

Show that the function $y = a \cos x + b \sin x$ is a solution to

$$\frac{d^2y}{dx^2} + y = 0$$

simply requires substituting in for y and $\frac{d^2y}{dx^2}$ in the left-hand side and checking the result is zero.

Sometimes questions are more straight-forward than people realise. For example,

Show that the function $y = a \cos x + b \sin x$ is a solution to

$$\frac{d^2y}{dx^2} + y = 0$$

simply requires substituting in for y and $\frac{d^2y}{dx^2}$ in the left-hand side and checking the result is zero.

Some people waste time solving the equation from scratch.

Sensible answers

Often there will be a way to think about whether the answer you have got is sensible.

Often there will be a way to think about whether the answer you have got is sensible. Exam markers despair when they see answers which are not sensible at all.

Often there will be a way to think about whether the answer you have got is sensible. Exam markers despair when they see answers which are not sensible at all.

- if you've taken the scalar product of two vectors, do you have a scalar?

Often there will be a way to think about whether the answer you have got is sensible. Exam markers despair when they see answers which are not sensible at all.

- if you've taken the scalar product of two vectors, do you have a scalar?
- If you have found an indefinite integral, does it differentiate to what you expected?

Often there will be a way to think about whether the answer you have got is sensible. Exam markers despair when they see answers which are not sensible at all.

- if you've taken the scalar product of two vectors, do you have a scalar?
- If you have found an indefinite integral, does it differentiate to what you expected?
- If you have found the maximum point on a function, does it look like a maximum?

Often there will be a way to think about whether the answer you have got is sensible. Exam markers despair when they see answers which are not sensible at all.

- if you've taken the scalar product of two vectors, do you have a scalar?
- If you have found an indefinite integral, does it differentiate to what you expected?
- If you have found the maximum point on a function, does it look like a maximum?

Often simple questions like this can help to spot errors and pick up extra marks.

Checking at the end

Often at the end of the exam, the last thing you want to do is look back at your answers.

Often at the end of the exam, the last thing you want to do is look back at your answers. But it's quite likely you'll be able to find an error or two if you do so.

Often at the end of the exam, the last thing you want to do is look back at your answers. But it's quite likely you'll be able to find an error or two if you do so.

If you have time left over, it's a really good idea to check through some of your algebra.

Often at the end of the exam, the last thing you want to do is look back at your answers. But it's quite likely you'll be able to find an error or two if you do so.

If you have time left over, it's a really good idea to check through some of your algebra. An extra few marks can move you up a grade, or stop you from failing.

Still to come

The final videos should go out on Thursday of Week 10.

The final videos should go out on Thursday of Week 10.
There will be two classes in Week 11 as usual, and no classes in Week 12.

The final videos should go out on Thursday of Week 10. There will be two classes in Week 11 as usual, and no classes in Week 12.

The exam period starts straight after Week 12, so not much time for revision. A revision tutorial sheet will be available on the module website in Week 11.

The final videos should go out on Thursday of Week 10. There will be two classes in Week 11 as usual, and no classes in Week 12.

The exam period starts straight after Week 12, so not much time for revision. A revision tutorial sheet will be available on the module website in Week 11.

There will be two drop-in sessions:

- Tuesday, 14th May: 2–5pm (Diamond WR3)
- Thursday, 16th May: 1–4pm (Diamond WR3)

We hope you enjoy the rest of the course!