## PARTIAL SECOND DERIVATIVES

**Announcement.** Remind students that there is a full-class lecture in Week 8 (MAS140 W1 SU AUD, MAS151 W1 DIA LT4, MAS152 M2 Church, MAS156(Elec) M5 DIA LT4, MAS156(Aero) Tu1 DIA LT1, MAS161 M5 DIA LT4). All should attend.

**5 minute review.** Remind students how to calculate the second derivatives of f(x,y), the notation  $f_{xx}$ ,  $f_{yy}$  etc, and that  $f_{xy} = f_{yx}$  (a least in every case they'll come across). Remind them how to find stationary points  $(f_x = f_y = 0)$ , and that to test them look at  $\Delta = f_{xx}f_{yy} - (f_{xy})^2$ , with

- $\Delta > 0$  and  $f_{xx} > 0$  giving a local minimum;
- $\Delta > 0$  and  $f_{xx} < 0$  giving a local maximum;
- $\Delta < 0$  giving a saddle;
- $\Delta = 0$  being inconclusive.

Class warm-up. Find the stationary points of  $f(x,y) = 2x^3 + 6xy^2 - 3y^3 - 150x$ . Emphasise how good presentational logic is needed to make sure all four stationary points are found. Perhaps leave students to classify the points.

**Problems.** Choose from the below.

- 1. Stationary points. Find and classify the stationary points of
  - (a)  $f(x,y) = 3x^2 + xy + 2y$ ;
  - (b)  $f(x,y) = e^{-(x^2+y^2)}$ ;
  - (c)  $f(x,y) = \cos(xy)$ . (Hint: try to think about what the surface z = f(x,y) looks like, investigating well-chosen contours.)
- 2. Harmonic functions. Let  $f(x,y) = e^{-y} \cos x$ .
  - (a) Show that  $\frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} = 0$ . (Functions satisfying this equation are called *harmonic* functions.)
  - (b) Does this function have any stationary points?
  - (c) What do slices through the surface z = f(x, y) parallel to the x-axis look like (e.g. where y = 0, y = 1 etc)? Can you describe the surface overall?
- 3. **Taylor series**. Just as in the single variable case, it is possible to approximate well-behaved functions of two variables with a *Taylor series* at (0,0):

$$f(x,y) = f(0,0) + xf_x(0,0) + yf_y(0,0)$$
  
+  $\frac{1}{2!} \left( x^2 f_{xx}(0,0) + 2xy f_{xy}(0,0) + y^2 f_{yy}(0,0) \right) + \dots$ 

for small x and y.

- (a) Find the first few terms in the Taylor series of  $f(x,y) = e^x \ln(1+y)$ . Try some points (x,y) near (0,0) to see how well your approximation works.
- (b) Can you guess how the Taylor series for f(x,y) continues past the '...'?

1

For the warm-up, there are four stationary points, namely a minimum at (5,0), a maximum at (-5,0), and saddles at (3,4) and (-3,-4).

## Selected answers and hints.

- 1. (a) There is a saddle at (-2, 12).
  - (b) There is a maximum at (0,0).
  - (c) There are maxima whenever xy is an even multiple of  $\pi$ , and minima whenever xy is an odd multiple of  $\pi$ . ( $\Delta = 0$  for the stationary points here, so thinking about how the function behaves is necessary.)
- 2. (b) There are no stationary points on this surface.
  - (c) Each slice through the surface parallel to the x-axis is a cosine wave, and moving in the positive y-direction, the amplitude of the wave decreases.
- 3. (a)  $e^x \ln(1+y) \approx y + xy \frac{y^2}{2} + \dots$ 
  - (b) The next term is

$$\frac{1}{3!} \left( x^3 f_{xxx}(0,0) + 3x^2 y f_{xxy}(0,0) + 3xy^2 f_{xyy}(0,0) + y^3 f_{yyy}(0,0) \right).$$

For more details, start a thread on the discussion board.