

## THE ARGAND DIAGRAM

**5 minute review.** Review the argand diagram (aka argand plane, aka complex plane), the modulus and argument and the polar form of a complex number as  $z = r(\cos \theta + i \sin \theta)$ . Also cover the geometric effect of addition and multiplication.

**Class warm-up.** With the help of students, calculate  $i^n$  for  $n = 1, \dots, 6$ , then plot them on the argand plane. Do the same for  $n = -1, -2$ , etc. Next plot  $(1+i)^n$  for  $n = 1, \dots, 6$ . Then plot  $(1+i)^0$  and ask students to guess where  $(1+i)^{-1}$  and  $(1+i)^{-2}$  sit. Confirm with calculations.

**Problems.** Choose from the below.

1. **Plotting complex numbers.** Calculate the modulus and argument of the following, plotting them on the argand diagram.

$$(a) \sqrt{3} + i, (b) -\sqrt{2} + \sqrt{6}i, (c) \frac{1}{2 + 3i}.$$

2. **Loci.**

- (a) What does the set  $\{z \in \mathbb{C} : |z| = 3\}$  look like in the argand plane (that is, those points in the complex plane for which  $|z| = 3$ )?
- (b) For  $z$  and  $w$  complex numbers, plot  $z$ ,  $w$ ,  $-w$  and  $z - w$  on the argand plane. What is the geometric meaning of  $|z - w|$ ? Which complex numbers  $z$  satisfy  $|z - i| = |z - 1|$ ? Draw them on the complex plane.

3. **Sums of roots.**

- (a) Which complex number has modulus 1 and argument  $\frac{2\pi}{5}$ ? What is  $z^5$ ? Plot  $z$ ,  $z^2$ ,  $z^3$ ,  $z^4$  and  $z^5$  on the argand diagram. Without calculating, what is  $z^4 + z^3 + z^2 + z$ ? (Hint:  $z^5 - 1 = 0$ , but  $z \neq 1$ . Factorise!)
- (b) If  $|z| = 2$  and  $\arg z = \frac{\pi}{3}$ , what is  $z^5 + 2z^4 + 4z^3 + 8z^2 + 16z$ ?

4. **Machin's Formula\*.**

- (a) Use the Binomial Theorem to write  $(5+i)^4$  in the form  $a+bi$ . Now show

$$(5+i)^4(239-i) = 114244(1+i).$$

Take arguments of both sides to show


$$4 \arctan(1/5) - \arctan(1/239) = \pi/4. \quad (*)$$

- (b) The Maclaurin expansion for  $\arctan x$ , valid for  $|x| \leq 1$ , is

$$\arctan x = x - \frac{1}{3}x^3 + \frac{1}{5}x^5 - \dots$$

Use these three terms to calculate (to 8 d.p.) an approximation to the left hand side of (\*). Does it look like  $\pi$  after multiplying by 4?


For the warm-up, here is a diagram:



### Selected answers and hints.

- $\sqrt{3} + i$  has modulus 2 and argument  $\frac{\pi}{6}$ .
  - $-\sqrt{2} + \sqrt{6}i$  has modulus  $2\sqrt{2}$  and argument  $\frac{2\pi}{3}$ .
  - $\frac{1}{2+3i} = \frac{2-3i}{2^2+3^2} = \frac{2}{13} - \frac{3}{13}i$ , which has modulus  $\frac{1}{\sqrt{13}}$  and argument  $-\tan^{-1}(\frac{3}{2}) \approx -0.98$  (2 s.f.).
- $\{z \in \mathbb{C} : |z| = 3\}$  is a circle, centred on the origin, radius 3.
  - $|z - w|$  gives the distance between  $z$  and  $w$ .

$\{z \in \mathbb{C} : |z - i| = |z - 1|\}$  is the set of all points whose distance from  $i$  is the same as their distance from 1 in the argand diagram. In other words, it is the perpendicular bisector of those points, namely the diagonal  $\text{Re}(z) = \text{Im}(z)$  (i.e.  $y = x$ ).



- Here,  $z = \cos(\frac{2\pi}{5}) + i \sin(\frac{2\pi}{5})$ , which is at a distance 1 from the origin, and a fifth of a full turn anticlockwise from the real axis. It follows that  $z^5$  will also have modulus 1 and argument  $5 \times \frac{2\pi}{5} = 2\pi$ ; that is,  $z^5 = 1$ .  
  
For the second part, since  $z^5 = 1$ , it follows that  $z^5 - 1 = 0$ , which factorises to give  $(z - 1)(z^4 + z^3 + z^2 + z + 1) = 0$ . Since  $z \neq 1$ , that means that  $z^4 + z^3 + z^2 + z = -1$ .
  - Here,  $z^6$  will have modulus  $2^6 = 64$  and argument  $6 \times \frac{\pi}{3} = 2\pi$ , so  $z^6 = 64$ . Hence,  $z^6 - 64 = 0$ , which factorises as  $(z - 2)(z^5 + 2z^4 + 4z^3 + 8z^2 + 16z + 32) = 0$ . It follows that  $z^5 + 2z^4 + 4z^3 + 8z^2 + 16z = -32$ .

For more details, start a thread on the discussion board.